



Fumagri® OPP

Food safety begins
in your poultry sheds

THE RESPONSIBLE CHOICE FOR FIGHTING SALMONELLA


PRODUCT

- Bactericidal and fungicidal disinfectant with OrthoPhenylPhenol (OPP)
- Dispersion by ULTRADIFFUSION®
- Complies with biocide regulations, use PTO3: livestock premises disinfection

ADVANTAGES

- Demonstrated effectiveness against salmonella, aspergillus and other bacteria, and moulds as well as against following viruses: *Avian influenza virus (H9N2)*, *Newcastle disease virus*, *Infectious bursal disease virus*
- Uniform diffusion of the disinfectant
- Safe to operators and animals: no humans or animals presence during treatment
- Convenient and fast application

RESULTS



Required efficacy according to standard NFT 72281: 5 log for bacteria and 4 log for moulds.

Virus: efficacy threshold: 4 log; Study of Institute of Animal Medicine, College of Veterinary Medicine, Gyeongsang National University, Chinju, Republic of Korea. Director: Hu-Jang Lee, DVM, PhD.




Fumagri® OPP

*Food safety begins
in your poultry sheds*

APPLICATION : Fumagri® OPP 1250 m³

USE To prevent and eradicate fungal and bacterial contamination
Optimal application conditions:
Minimum temperature: 9°C
Humidity 50-80%



Type of shed	Number of Fumagri® OPP 1250m ³ tins
400m ²	1
800m ²	2
1000m ²	3
1200m ²	4 (3 for side access shed)
Laying premises	To suit shed volume

Remember to disinfect the silo with Fumagri® OPP 20g
2 days before feed loading,
Dry treatment, no humidity input.

Use PTO4: storage of livestock feed.



Also available for other applications in 20g, 40g, 120g, 200g, 400g, 1000g pre-dosed tins.

Fumagri® OPP: biocidal use PTO3 and PTO4. Composition: OrthoPhenylPhenol, non-corrosive.
Take care when using biocides. Read label and product information before using.
Technical data sheet on request
design & layout: eretic brand consulting & design - photo credits: ©iStockphoto.com/barbara gibbons